

APPENDIX I

# Genes VII

Benjamin Lewin

(BEST) AVAILABLE COPY

OXFORD  
UNIVERSITY PRESS

OXFORD

UNIVERSITY PRESS

Great Clarendon Street, Oxford OX2 6DP

Oxford University Press is a department of the University of Oxford.  
It furthers the University's objective of excellence in research, scholarship,  
and education by publishing worldwide in  
Oxford New York

Athens Auckland Bangkok Bogotá Buenos Aires Calcutta  
Cape Town Chennai Dar es Salaam Delhi Florence Hong Kong Istanbul  
Karachi Kuala Lumpur Madrid Melbourne Mexico City Mumbai  
Nairobi Paris São Paulo Singapore Taipei Tokyo Toronto Warsaw  
with associated companies in Berlin Ibadan

Oxford is a registered trade mark of Oxford University Press  
in the UK and in certain other countries

Published in the United States  
by Oxford University Press Inc., New York

© Oxford University Press and Cell Press, 2000

The moral rights of the author have been asserted  
Database right Oxford University Press (maker)

First published 2000

All rights reserved. No part of this publication may be reproduced,  
stored in a retrieval system, or transmitted, in any form or by any means,  
without the prior permission in writing of Oxford University Press,  
or as expressly permitted by law, or under terms agreed with the appropriate  
reprographics rights organization. Enquiries concerning reproduction  
outside the scope of the above should be sent to the Rights Department,  
Oxford University Press, at the address above

You must not circulate this book in any other binding or cover  
and you must impose this same condition on any acquirer

A catalogue record for this book is available from the British Library

Library of Congress Cataloging in Publication Data  
(Data applied for)

ISBN 0-19-879276-X (Hbk)

Typeset by J&L Composition Ltd, Filey, North Yorkshire  
Printed in The United States of America

MASS. INST. TECH.

JAN 4 2002

LIBRARIES

BEST AVAILABLE COPY

QH430

L487


2000

operator, and is therefore constitutive like the *lacI<sup>-</sup>* alleles. Because the *lacI<sup>-</sup>* type of mutation inactivates the repressor, it is recessive to the wild type. However, the *-d* notation indicates that this variant of the negative type is dominant when paired with a wild-type allele. Such mutations are said to be *trans*-dominant; they are also called dominant negatives.


The reason for the dominance is that the *lacI<sup>-d</sup>* allele produces a "bad" subunit, which is not only itself unable to bind to operator DNA, but is also able as part of a tetramer to prevent any "good" subunits from binding. This demonstrates that the repressor tetramer as a whole, rather than the individual monomer, is needed to achieve repression. The poisoning effect also can be

**Figure 10.8**

Mutations that inactivate the *lacI* gene cause the operon to be constitutively expressed, because the mutant repressor protein cannot bind to the operator.



**Figure 10.9** Mutations map the regions of the *lacI* gene responsible for different functions. The DNA-binding domain is identified by *lacI<sup>-d</sup>* mutations at the N-terminal region; *lacI<sup>-</sup>* mutations unable to form tetramers are located between residues 220–280; other *lacI<sup>-</sup>* mutations occur throughout the gene; *lacI<sup>s</sup>* mutations occur in regularly spaced clusters between residues 62–300.



BEST AVAILABLE COPY

The diagram illustrates the cell cycle and its regulation. The cell cycle is shown as a circular process with phases G0, G1, S phase, G2, and M. Key components and regulators include:

- G0 Phase:** Cells can enter G0 from G1. Cyclin D and Cdk4,6 are associated with this phase.
- G1 Phase:** Contains the E2F transcription factor and the RB protein. Cyclin E and Cdk2 are associated with this phase.
- S Phase:** DNA replication occurs. Phosphorylation (P) is indicated.
- G2 Phase:** Contains cyclin A and B, and Cdc2.
- M Phase:** Mitosis.
- Tumor Suppressors:** p16, p27, p21, and p53 are shown. p16 inhibits Cdk4,6. p27, p21, and p53 are part of a regulatory network that inhibits the G1 to S transition.

But all the transforming forms of p53 turned out to be mutant forms of the protein! They fall into the category of dominant negative mutants, which function by overwhelming the wild-type protein and preventing it from functioning. The most common form of a dominant negative mutant is one that forms a heteromeric

Figure 28.24 shows that the same phenotype is produced either by the deletion of both alleles or by a missense point mutation in one allele that produces a dominant negative subunit. Both situations are found in human cancers. Mutations in p53 accumulate in many types of human cancer, probably because loss of p53 provides a growth advantage to cells; that is, wild-type p53 restrains growth. The diversity of these cancers suggests that p53 is not involved in a tissue-specific event, but in some general and rather common control of cell proliferation; and the loss of this control may be a secondary event that occurs to assist the growth of many tumors. Mutant p53 cells also have an increased